

A fungicide for the control of late potato blight. FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE FOR PROFESSIONAL USE ONLY

•

vater dispersible granule (WG) for GROUP U27 FUNGICIDE

staining 45% w/w cymoxanil.

PROTECT FROM FROST

 $ymbal_{45}$

IMPORTANT INFORMATION
For use only as an agricultural fungicide Maximum number of applications Maximum total dose 2.5 kg/ha 7 days before

Control of late blight of potatoes caused by Phytophthora infestant is achieved when used in containing active substances with multi-site activity

READ THE LABIL BEFORE USE. USING THIS PRODUCT HA MANKET HATTS INCONSISTENT WITH THE LABIL MAY BE AN OFEINGE. FOLLOW THE PRINCIPLES OF GOOD PLANT PROTECTION PRACTICE FOR USING PLANT PROTECTION AND OFFICE FOR USING PLANT PROTECTION.

KAND SAFETY INFORMATION

y cause an allory)c kin reaction
user as of see a further or
user as of see and of see and
special of demands of the under child
special of demands the unborn child
see user demands the unborn child
post cause demands in the unborn child
special of the unborn of the unborn child
special of the unborn child
sp ning 45% w/w cymoxanil

stective gloves
rd or concerned. Get medical advice/attention
sillage.

avoid risks to human health and the environment, comply with the instructions of contents/container to a licerteed hazavdous-waste disposa, contractor or nate except for empty clean triple mixed containers which can be disposed rezardous waste.

Whelehan end protection

BELCHIM

etion Limited doeshire PE19 8EW | Tel: 0044:0) 1480 403333 | Fax: 0044 (0

SAFETY PRECAUTIONS

Operator protection

Engineering control of operator exposure must be used where in Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal operative engineering control of the protective engineering expected protective engineering expected when handling surfake PROTECTIVE COLOVES when handling to the protection of the pr

contaminated surfaces.
WASH SPLASHES from skin immediately.
WASH HANDS AND EXPOSED SKIN before eating and crinking or

smoking and after work.
WHEN USING DO NOT EAT, DRINK OR SMOKE.
AVOID CONTACT WITH SKIN.

Environmental protection

DO NOT CONTAMINATE WATER with the product or its container (Do not clean application equipment near surface water (Avoid constitutions not drains from farmywist, and coast proposate containment to avoid environmental containment.)

Stonge and disposal

Stonge and disposal

KEEP IN ORIGINAL CONTANER (to may closed, in a safe place
EMPTY CONTINUES COMPLETELY and dispose of carlely.

KEEP OUT OF BEACH OF CHILDREN

KEEP AWAY FROM FOOD, DAINK AND ANIMAL

FEEDINGSTUFFS.

FIRST MAY FROM FOOD, DAINK AND ANIMAL

FEEDINGSTUFFS.

RESTRICTIONS
Allow a minimum of 7 days between applications. Only apply
CYMBAL* 35 in tank minimum with other fungicides containing
additinent active ingredient. The container must not be re-used

RESSTANCE
When functions with the same mode of action are used
When functions exercis years in the same field, naturally
occurring less seneither stains may autowne, propagate and
become sominant in that field, it is therefore advised that
where the probles proxy programmes should include fungibiles
with different modes of action.

DIRECTIONS FOR USE IMPORTANT: This information is approved as part of the IMPORTANT: This informations within this section must be read carefully in order to obtain safe and successful use of this

CROPS
CYMBAL® 45 can be used on all varieties of early and main crop potatoes. On crops grown for processing, consult your processor before use.

TIMING AND WEATHER

Timing of spray applications: should take into account the normal a prevalence of the disease in the area and concilions in the particular season. The first spray should be applied as soon as there is a risk of blight infection or an official warming has been instants, whichever occurs first. In the absence of a warming, spraying should begin just before the crop meets along the rows.

DOSE

CYMBAL® 45 should be applied at 0.25 kg/ha in tank-mixture with one of the following products at the recommended rate of application:

Fluazinam (500 g/L) e.g. Shirlan. This mixture will give a in tuber blight.

Cyazofamid e.g. Ranman Top. This mixture will give a reduction tuber blight.

All other requirements and restrictions on the partner product label should be adhered to.

VOLUME AND APPLICATION

Apply CYMBAL® 45 in not less than 200 litres of water per hectare using a MEDIUM quality spray (as defined by BCPC).

MIXING

Spraying equipment should be thoroughly cleaned before spraying CYMBAL 45 mixer solly with votor and should be added directly to the spray tank during filling. Wash out container and add washings to the spray tank. Continue agitation until spraying is completed.

NOTICE TO BUYER

We would be supplied by us are of high glade and we believe them all goods supplied by us are of high glade and we believe them to be suitable for any purpose for which we expressly supply them, but as we cannot exercise control over their mixing or use, e all conditions and warnnities statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded and in one tepponibility will be accepted by ut for any stamper of injury whatscever arising from their storage, handling, application or vivo.

in WARRANTY

WE Shausive research has pointed out that the product, if applied or Eshausive research has pointed out that the product, if applied or according to the distributor's prescriptions, is suitable for the given usages, Since its concervation, handling, transportation at and possible misuse are beyond the distributor will deny any the responsibility as to the results and/or any damage that might eshausive from the product's concervation, handling, transportation on possible misuse, the visil only guarantee a contract quality of the product, in case of doubt about or unclarity of the usage the product, in case of doubt about or unclarity of the usage free precipions for men and/or undernibur cultures or races, please contact the distributor for more information.

Applications should be repeated at 10-4d by internals depending on the propilers colline it considers in indexion. A 7 day puty internal may be used under exceptional at the summanders e.g., slight test at evaluable high and is forecast, at the summanders e.g., slight test at evaluable high and is forecast, at the summanders e.g., slight test at evaluable high and is forecast, at the summanders e.g., slight test at evaluable high and one, allows of the summanders e.g., slight test at evaluable high test at the summanders e.g., slight test at the slight test at the summanders e.g., slight test at the slight test reduction



www.belchim.co.uk/pdf/MSDS/ Cymbal_MSDS.pdf