

A fungicide for the control of late foliar blight (Phytophthora infestans).

FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE

MAPP No.: 16750

A water dispersible granule (WG) formulation containing 45% w/w cymoxanil.

	GROUP	U27 FUNGICIDE	
		TANT INFORMATION	GICIDE
Crop	Maximum individual dos (kg product/ha)	e Maximum number of treatments (per year)	Latest time of application
Potato	0.25	10	7 days before harvest
Do not re-t	ific restrictions: interval of 7 days must be obs- use container for any purpose. IS CABEL BEFORE USE. USING IE LABEL MAY BE AN OFFENCE PRO	THIS PRODUCT IN A MANNE	R THAT IS INCONSISTENT CTICE FOR USING PLANT



H361fd: Suspected of damaging feetility Suspected of demaying the unborn child H373 : May cause damage to organs through prolonged or repeated exposure (blood, thymus) H410: Very texic to aquatic life with long lasting effects

FS+ID: Do not breatly sayye TOTAL August release who she area increases and Pale Wew protective glans

1638 + P313 'F experieu er cur curreu L'et medical advice/ attention. #391: Collect spillade


P501: Dispose of contents/container to a licensed hazardour-waste disputal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

EUH401: To avoid risks to human health and the environment, comply with the instructions for use.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.



UFI: 4HH8-J045-J00M-VR9X



SAFETY PRECAUTIONS

Operator protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

- WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling the product.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

- WASH SPLASHES from skin immediately.

- WASH HANDS AND EXPOSED SKIN before eating and drinking or smoking and after work.

- WHEN USING DO NOT EAT, DRINK OR SMOKE. - AVOID CONTACT WITH SKIN

- WEAR SUITABLE GLOVES.

Environmental protection-

- DO NOT CONTAMINATE SURFACE WATERS OR DITCHES with Chemical or used container.

- Use appropriate containment to avoid environmental contamination.

Storage and disposal

- KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place - EMPTY CONTAINER COMPLETELY and dispose of safely.

- KEEP OUT OF REACH OF CHILDREN

- KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDINGSTUFFS - THIS MATERIAL AND ITS CONTAINER must be disposed of in a

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

RESTRICTIONS

flow a minimum of 7 days between applications.

Only apply DRUM* in tank mixture with other fungicides containing different active ingredient.

RESISTANCE MANAGEMENT

When fungicides with the same mode of action are used repeatedly over several years in the same field, naturally occurring less sensitive strains may survive, propagate and become dominant in that field. It is therefore advised to design spray programmes to utilise fungicides with different modes of action and observe manufacturers recommendations on dose, timing and spray interval. Cymoxanii must always be used in mixture with actives of a different mode of action and alternate in the spray programme with other mixed formulations of active ingredients with different modes of action or fungicide families.

CROPS

DRUM® can be used on all varieties of early and main crop potatoes. On crops grown for processing, consult your processor before use. TIMING AND WEATHER

Timing of spray applications should take into account the normal prevalence of the disease in the area and conditions in the particular season. The first spray should be applied as soon as there is a risk of blight infection or an official warning has been issued, whichever occurs first.

In the absence of a warning, spraying should begin just before the

crop meets along the rows.

Applications should be repeated at 10-14 day intervals depending on the prevalence of the disease and favourable conditions for infection. A 7 day spray interval may be used under exceptional circumstances, e.g. blight risk is already high and rain is forecast at the time the next 10 day spray is due. High risk situations include conditions of rapid growth, irrigated crops, areas of Intensive potato growing, particularly blight sensitive varieties, when further warnings are issued, or when active blight is in the area. The longer Interval is recommended under dry conditions, Irrigation of greet increases the risk of blight and crops should always be sprayed with DRUM® as soon as possible after impation subject to the minimum spray interval. Irrigated crops should normally be sprayed at 10 day

DOSE

DRUM® should be applied at 0.2-0.25 kg/ha in tank-mixture with one of the following products: the highest rate of DRUM® should be used when conditions are lavourable for P. Infestans-and disease pressure is high

Fluazinam (Shirlan, MAPP 18406) at 0.3-0.4 litres product per hectare (rate of Shirlan is dependent on blight risk and spraynterval, see product labell. This mixture will give a reduction in

Cyazofamid (Ramman Top MAPP 14753) at 500 ml of product per hectare. This mixture will reduce the percentage of blight affected tubers due to the activity of Ranman Top.

All other requirements and restrictions on the partner product label should be adhered to.

VOLUME AND APPLICATION

Apply DRUM* in not less than 200 litres of water per hectare using a MEDIUM quality spray (as defined by BCPC).

Spraying equipment should be thoroughly cleaned before spraying. DRUM® mixes easily with water and should be added directly to the spray tank during filling. Wash out container and add washings to the spray tank. Continue agitation until spraying is complete.

NOTICE TO BUYER

All goods supplied by us are of high grade and we believe them to be suitable for any purpose for which we expressly supply them, but as we cannot exercise control over their mixing or use, all conditions and warranties statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use.

Exhaustive research has pointed out that the product, if applied according to the distributors' prescriptions, is suitable for the given usages. Since its conservation, handling, transportation and possible misuse are beyond the distributors' control - as are unforeseen circumstances, the distributor will deny any responsibility as to the results and/or any damage that might ensue from the products' conservation, handling, transportation or possible misuse. He will only guarantee a constant quality of the product. In case of doubt about or unclarity of the usage prescriptions for new and/ or unfamiliar cultures or races, please contact the distributor for more information.

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:



www.belchim.co.uk/pdf/MSDS/Drum MSDS.pdf

Marketing Company and approval holder Belchim Crop Protection Limited 1b Fenice Court Phoenix Park | Eaton Socon, St. Neots | Cambridgeshire PE19 8EW Tel.: 01480 405535 | Fax: 01480 405444 | info-uk@belchim.com | www.belchim.co.uk