

FUNGICIDE



UK

Content: **5**L e

A fungicide for use on apple and pear against *Botrytis spp.* and *Gloeosporium spp.* storage diseases

A suspension concentrate containing 336 g/L pyrimethanil (30.59 % w/w) and 133 g/L fludioxonil (12.11 % w/w).

MAPP No.: 18244

GROUP	9	12	FUNGICIDES
-------	---	----	------------

Production date / Batch number: see packaging

**PROTECT FROM FROST
FOR PROFESSIONAL USE ONLY**




The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.



© Pomax is a registered trademark of Globachem NV.

For warehouse purpose only:



Distributed by: **Belchim Crop Protection Limited**
1b Fenice Court, Phoenix Park | Eaton Socon, St. Neots | Cambridgeshire PE19 8EW
Tel.: 01480 403333 | Fax: 01480 403444 | Info-uk@belchim.com | www.belchim.co.uk

APPROVAL HOLDER: **GLOBACHEM NV**
Lichtenberglaan 2019, B-3800 Sint-Truiden
www.globachem.com



SAFETY PRECAUTIONS

OPERATOR PROTECTION

Engineering control of operator exposure must be used where reasonably practicable in addition to the following:

- WEAR SUITABLE PROTECTIVE GLOVES when handling the concentrate.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows that they provide an equal or higher standard of protection.

ENVIRONMENTAL PROTECTION

- Do not contaminate water with the product or its container (Do not clean application equipment near surface water/ Avoid contamination via drains from farmyards and roads).
- To protect aquatic organisms, respect an unsprayed buffer zone distance to surface water bodies in line with LERAP requirements.
- DO NOT ALLOW DIRECT SPRAY from broadcast air-assisted sprayers to fall within 20 m of the top of the bank of a static or flowing waterbody, unless a Local Environmental Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 5 m of the top of a ditch which is dry at the time of application. Aim spray away from water.
- This product qualifies for inclusion within the Local Environmental Risk Assessment for Pesticides (LERAP) Scheme. Before each spraying operation from a broadcast air assisted sprayer, either a LERAP must be carried out in accordance with CRD's published guidance or the statutory buffer zone must be maintained.
- The results of the LERAP must be recorded and kept available for three years.

STORAGE AND DISPOSAL

- DO NOT RE-USE CONTAINER for any purpose.
- KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.
- KEEP OUT OF REACH OF CHILDREN.
- KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.
- WASH OUT CONTAINER THOROUGHLY, empty washings into spray tank and dispose of safely.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

CROP SPECIFIC INFORMATION

POMAX can be used on apples and pears to control storage diseases (*Botrytis spp.* and *Gloeosporium spp.*). Two applications of 1.6 L/ha can be made starting from 6 weeks before harvest (from 4 months) with 7 day interval. Respect a pre-harvest interval of 5 days.

LEVELS OF CONTROL

Apples only: Control of *Gloeosporium spp.*

Pomax is **NOT** for the control of *Gloeosporium spp.* on pears.

Apples and pears: Reduction/some control of *Botrytis cinerea*.

RESTRICTIONS/ WARNINGS

Crop safety on Cox and Bramley varieties has not been established.

MIXING AND SPRAYING

Make sure the sprayer is set to give an even application at the correct volume. Fill the spray tank with half the required volume of clean water and start agitation. Add the required amount of POMAX and continue agitation whilst adding the rest of the water. Agitate the mixture thoroughly before use and continue agitation during spraying. Take particular care to avoid overlapping of spray swaths. Thoroughly wash all spray and measuring equipment with water immediately after use.

RESISTANCE MANAGEMENT

This product contains pyrimethanil which is a member of the anilino-pyrimidine (AP) fungicide group and fludioxonil which is a member of the phenylpyrrole (PP) fungicide group. Application should be made in accordance with the FRAC anilinopyrimidine (AP) working group *Botrytis* guidelines.

Where only 2 applications are made per season, the number of anilinopyrimidine products should be limited to 1. Where up to 6 *Botrytis* treatments are made per crop and season, a maximum of 2 applications of anilinopyrimidine-containing

products are recommended. Where more than 6 fungicide treatments are made per crop and season, a maximum of 3 applications of anilinopyrimidine-containing products is recommended.

In order to minimise the likelihood of the development of resistance, it is recommended that POMAX should be used in a programme with products of different chemical groups. Alternating or mixing fungicides with different modes of actions is a recognised disease resistance management strategy.

CONDITIONS OF SUPPLY

All goods supplied by us are of high grade and we believe them to be suitable but, as we cannot exercise control over their storage, handling, mixing or use, or the weather conditions before, during or after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded. No responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

**To access the Safety Data Sheet
for this product, scan the QR code
or use the weblink below:**



http://www.belchim.co.uk/pdf/MSDS/Pomax_MSDS.pdf

Alternatively, contact your supplier.

**IMPORTANT INFORMATION
FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE**

Crops	Max. individual dose: (L product / ha)	Maximum total dose: (L product / ha / crop)	Latest time of application
Apples and pears	1.6	3.2	5 days before harvest

Other specific restrictions:

Do not apply by hand-held equipment.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

Pomax®

A suspension concentrate containing 336 g/L pyrimethanil (30.59 % w/w) and 133 g/L fludioxonil (12.11 % w/w).

**WARNING**

H410 Very toxic to aquatic life with long lasting effects.

P280 Wear protective gloves

P391 Collect spillage

P501 Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

EUH401 To avoid risks to human health and the environment, comply with the instructions for use.

SP1 Do not contaminate water with the product or its container.

SPe3 To protect aquatic organisms respect a 20 m unsprayed buffer zone to surface water bodies.