



A fungicide for the control of late potato blight.

A water dispersible granule (WG) formulation containing 45% w/w cymoxanil.

IMPORTANT INFORMATION FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE

Crop	Maximum individual dose (kg product/ha)	Maximum number of applications	Maximum total dose	Latest time of application
Potatoes (seed and ware)	0.25	10 per crop	2.5 kg/ha	7 days before harvest

Specific diseases controlled: Control of late blight of potatoes caused by *Phytophthora infestans* is achieved when used in conjunction with products containing active substances with multi-site activity

Other specific restrictions:

- A minimum interval of 7 days must be observed between applications.
- Do not re-use container for any purpose.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE PRINCIPLES OF GOOD PLANT PROTECTION PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

RISK AND SAFETY INFORMATION

UFI: 4HH8-J045-J00M-VR9X

Cymbal® 45 - A water dispersible granule (WG) formulation containing 45% w/w cymoxanil

May cause an allergic skin reaction

Causes serious eye irritation

Suspected of damaging fertility

Suspected of damaging the unborn child

May cause damage to organs through prolonged or repeated exposure (blood, thymus)

Very toxic to aquatic life with long lasting effects

Do not breathe spray

Avoid release to the environment

Wear protective gloves

IF exposed or concerned: Get medical advice/attention

Collect spillage

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean triple rinsed containers which can be disposed of as non-hazardous waste

To avoid risks to human health and the environment, comply with the instructions for use

PCS N° 04647

GROUP 27 FUNGICIDE

FUNGICIDE

Contents:

1kg^e



PCS No.: 04647

Production date/batch n°:
see packaging

**FOR USE ONLY AS AN
AGRICULTURAL FUNGICIDE**

FOR PROFESSIONAL USE ONLY

PROTECT FROM FROST

**24 Hour emergency number
0032 14 58 45 45**

Distributed by:



Whelehan Crop Protection
Suite 12 Bunkilla Plaza
Bracetown Business Park
Clonee
Co. Meath
Ireland
Technical Enquiries: 01 4688900

Authorisation Holder :
Certis Belchim BV
Stadsplateau 16,
3521 AZ Utrecht,
PO Box 607, 3500 AP Utrecht,
Netherlands

® Registered trademark of
Belchim Crop Protection NV/SA

cymbal 45



SAFETY PRECAUTIONS

Operator protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

- WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling the product.
- WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.

WASH SPLASHES from skin immediately.

WASH HANDS AND EXPOSED SKIN before eating and drinking or smoking and after work.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

AVOID CONTACT WITH SKIN.

WEAR SUITABLE GLOVES.

Environmental protection

- DO NOT CONTAMINATE WATER with the product or its container (Do not clean application equipment near surface water /Avoid contamination via drains from farmyards and roads).
- Use appropriate containment to avoid environmental contamination.

Storage and disposal

- KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.
- EMPTY CONTAINER COMPLETELY and dispose of safely.
- KEEP OUT OF REACH OF CHILDREN.
- KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDINGSTUFFS.
- THIS MATERIAL AND ITS CONTAINER must be disposed of in a safe way.

RESTRICTIONS

Allow a minimum of 7 days between applications. Only apply **CYMBAL® 45** in tank mixture with other fungicides containing a different active ingredient. The container must not be re-used for any purpose.

RESISTANCE

When fungicides with the same mode of action are used repeatedly over several years in the same field, naturally occurring less sensitive strains may survive, propagate and become dominant in that field. It is therefore advised that wherever possible spray programmes should include fungicides with different modes of action.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

CROPS

CYMBAL® 45 can be used on all varieties of early and main crop potatoes. On crops grown for processing, consult your processor before use.

TIMING AND WEATHER

Timing of spray applications should take into account the normal prevalence of the disease in the area and conditions in the particular season. The first spray should be applied as soon as there is a risk of blight infection or an official warning has been issued, whichever occurs first.

In the absence of a warning, spraying should begin just before the crop meets along the rows.

Applications should be repeated at 10-14 day intervals depending on the prevalence of the disease and favourable conditions for infection. A 7 day spray interval may be used under exceptional circumstances, e.g. blight risk is already high and rain is forecast at the time the next 10 day spray is due. High risk situations include conditions of rapid growth, irrigated crops, areas of intensive potato growing, particularly blight sensitive varieties, when further warnings are issued, or when active blight is in the area. The longer interval is recommended under dry conditions. Irrigation of crops increases the risk of blight and crops should always be sprayed with **CYMBAL® 45** as soon as possible after irrigation subject to the minimum spray interval. Irrigated crops should normally be sprayed at 10 day intervals. Maximum 10 applications.

DOSE

CYMBAL® 45 should be applied at 0.25 kg/ha in tank-mixture with one of the following products at the recommended rate of application:

Fluazinam (500 g/L) e.g. Shirlan. This mixture will give a reduction in tuber blight.

OR

Cyazofamid e.g. Ranman Top. This mixture will give a reduction in tuber blight.

All other requirements and restrictions on the partner product label should be adhered to.

VOLUME AND APPLICATION

Apply **CYMBAL® 45** in not less than 200 litres of water per hectare using a MEDIUM quality spray (as defined by BCPC).

MIXING

Spraying equipment should be thoroughly cleaned before spraying. **CYMBAL® 45** mixes easily with water and should be added directly to the spray tank during filling. Wash out container and add washings to the spray tank. Continue agitation until spraying is complete.

NOTICE TO BUYER

All goods supplied by us are of high grade and we believe them to be suitable for any purpose for which we expressly supply them, but as we cannot exercise control over their mixing or use, all conditions and warranties statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use.

WARRANTY

Exhaustive research has pointed out that the product, if applied according to the distributors' prescriptions, is suitable for the given usages. Since its conservation, handling, transportation and possible misuse are beyond the distributors' control - as are unforeseen circumstances, the distributor will deny any responsibility as to the results and/or any damage that might ensue from the products' conservation, handling, transportation or possible misuse. He will only guarantee a constant quality of the product. In case of doubt about or unclarity of the usage prescriptions for new and/or unfamiliar cultures or races, please contact the distributor for more information.

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:



WEBLINK to specific data sheet on manufacturer's website:

www.certisbelchim.co.uk/pdf/MSDS/Cymbal_IE_MSDS.pdf

Alternatively, contact your supplier.



Marketing company: **Certis Belchim BV**
Suite 5, 3 Riverside | Granta Park | Great Abington | Cambridgeshire | CB21 6AD
Tel.: +44 (0)1223 894261 | info.uk@certisbelchim.com | www.certisbelchim.com

UN 3077